

Blue and IR Light Curves of the Mysterious Pre-Main Sequence Star V582 Mon (KH 15D) from 1955 to 1970

Paolo Maffei¹, Stefano Ciprini^{1,2*}, and Gino Tosti¹

¹*Physics Department and Astronomical Observatory, University of Perugia, via Pascoli, 06123 Perugia, Italy*

²*Tuorla Astronomical Observatory, University of Turku, Väisäläntie 20, 21500 Piikkiö, Finland*

submitted to MNRAS

ABSTRACT

In recent years, an increasing number of publications have been addressed to the peculiar and mysterious pre-main sequence star V582 Mon, also known as KH 15D. This extraordinary T Tauri star, located in the young star cluster NGC 2264, appears as to be an eclipsing variable. In the present paper, we report a unique and self-consistent set of light curves in the blue and near-infrared bands, spanning a 15-year interval (epoch 1955–1970). Our photometric data show clearly the beginning of the eclipse stage occurred in early 1958 in the blue, and perhaps around four years later in the infrared. The light curve period turns out to be the same reported by recent observations (about 48.3 days), so that no evidence for a period change results. On the other hand, in our data the light curve shape appears as sinusoidal and is therefore different from the one displayed today. The photometric behaviour, determined with time-series and colour-index analysis, suggests that V582 Mon (KH 15D) could be initially surrounded by an accretion disk/torus seen edge-on, with subsequent thin dust formation at the beginning of the blue radiation absorption. The dust could then aggregate into larger particles providing the transition between selective and total absorption, accompanied with eclipsing variability in the infrared. The minima of the periodic light curve become deeper due to the increasing dimension and number of dust grains, and then flattens due to a contraction in the disk.

Key words: stars: individual (V582 Mon) — stars: individual (KH 15D) — stars: variables: other — stars: pre-main sequence — stars: circumstellar matter — techniques: photometric — stars: planetary systems: protoplanetary disks

1 INTRODUCTION

The variable star V582 Mon ($\alpha = 6^{\text{h}} 41^{\text{m}} 10.3^{\text{s}}$, $\delta = +09^{\circ} 28' 34''$ (J2000.0)) was discovered by Badalian & Erastova (1972). It is classified as a type INS (i.e. Orion variable with rapid light changes: up to 1 mag in 1–10 days) in the GCVS (Kholopov 1985). This star was re-discovered as an eclipse variable by Kearns & Herbst (1998) during a systematic study of young star variability in the region of the cluster NGC 2264. The star, named 15D in a list of 209 objects (Kearns et al. 1997), appeared as an eclipsing variable with an amplitude of about 3 mag in the I_c (Cousins) band, with a period of 48–49 days. This was a unique case among the very young stars observed as discussed by Kearns et al. (1997). The star is also designated as n. 150 in the list of Park et al. (2000) and as n. 391 in the list of Flaccomio et al. (1999).

Table 1. The number of Blue and IR photographic plates in the field of V582 Mon (KH 15D) from 1955 to 1970.

Telescope	N Blue	Years	N IR	Years
Reflector Loiano 0.60m	70	1957-59	9	1957-59
Reflector Asiago 1.22m	14	1955-63	5	1956-63
Schmidt Asiago 0.50/0.40m	52	1959-62	38	1959-65
Schmidt Asiago 0.90/0.65m	42	1967-70	47	1967-70
Schmidt Bergedorf 1.20/0.80m	5	1962-64	...	...

The depth (3.5 mag) and the long duration of the eclipse (about 40% of each cycle, which lasts for 48.4 days) rule out a companion star as the cause of this event (Hamilton et al. 2001; Herbst et al. 2002). Polarization levels of 2% observed at half eclipse and 0.2 ± 0.2 observed out of the eclipse (Agol et al. 2004), would imply that the obscuring material is composed of rather large particles. In order to explain these phenomena, two hypotheses were formulated: 1) a dou-

* offprints: stefano.ciprini@pg.infn.it




Figure 1. The complete light curves in the blue ($\sim B$ band) and photographic IR ($\sim I_c$ band) of V582 Mon (KH 15D) from 1955 to 1970 (due to the large plot scale nearby observations appear as a single point).

ble star obscured in different ways by circumstellar matter; 2) a single star surrounded by a peculiarly warped or ridged proto-planetary disk. In the second case, the light of the star is periodically intercepted by this dusty ridge orbiting around the star in a plane placed edge-on with respect to the line of sight.

Actually, there is a large consensus that the eclipses are caused by circumstellar material, but it is not clear what is the composition and the spatial distribution of this material. Theories include an edge-on circumstellar disk with a mass density wave or a disk deformation induced by a proto-planet (Hamilton et al. 2001; Herbst et al. 2002; Winn et al. 2003; Agol et al. 2004), an orbiting vortex of solid particles (Barge & Viton 2003), and an asymmetric common envelope (Grinin & Tambovtseva 2002). V582 Mon (KH 15D) is quite unique and extremely interesting because the fortuitous alignment may allow us to study the circumstellar (or even protoplanetary) processes, and because the eclipse provides a natural “coronagraph” that can be used to map out the environment of the underlying T Tauri star (Hamilton et al. 2003; Agol et al. 2004; Deming, Charbonneau, & Harrington 2004). In fact, this object should be a weak-line T Tauri star, still accreting matter from a circumstellar disk and endowed with a well-collimated bipolar jet (Hamilton et al. 2001). High resolution near-IR spectroscopy revealed the presence of prominent molecular hydrogen emission probably due to shock excitation of the ambient gas by a bipolar outflow from the circumstellar disk (Deming, Charbonneau, & Harrington 2004). As evidence of this hypothesis, there is also the observation of Tokunaga et al. (2004), who revealed clearly an H_2 emission filament next to the central star. Furthermore, spectro-polarimetric observations suggest that the star is completely eclipsed, so that the flux during

the eclipse is entirely due to scattered light. The scattering should be due to large dust grains ($\sim 10 \mu m$) similar to the interplanetary grains that scatter the zodiacal light (Agol et al. 2004). Recent high-resolution spectroscopic monitoring showed that V582 Mon is a single-line spectroscopic binary (Johnson et al. 2004). A model of gradual opaque-screen occultation was proposed by Winn et al. (2004), and a model with an eccentric binary that is encircled by a vertically thin, inclined ring of dust and gas was suggested by (Chiang & Murray-Clay 2004) in order to explain the available historical light curve prior to the present work.

The light curve of V582 Mon (KH 15D) from 1967 to 1982, based on the Asiago Observatory infrared plates, displays a periodic alternation between bright and faint states with a constant period of 48.35 days (Johnson & Winn 2004). An international observation campaign during 2001–2002 showed clearly a periodic variability recurring with a period of 48.36 days (Herbst et al. 2002). Observations during six seasons from 1995 to 2001 showed again a periodical variability like an eclipsing binary, with a period of 48.34 days (Hamilton et al. 2001). On the other hand, the analysis of the Harvard College Observatory archival plates from 1913 to 1951 showed that there was no variability, and the deep eclipses observed today did not occur in the first half of the 20th century (Winn et al. 2003). In that work, the need for further, more recent archival images was underlined, as they might have shown the onset of eclipses.

In our paper, we present the results of the analysis of the plate images obtained by one of us (PM) during 15 years in a systematic study of the variability of young stars in the cluster NGC 2264. The data presented and discussed here pertain to the star V582 Mon (KH 15D) and concern the observations performed in the blue (B) and near-infrared

Table 2. Our photometric measurements in the blue and IR of V582 Mon (KH 15D) in the period 1955-1970 (for the meaning of the telescope acronyms see Table 1).

Date [JD-2430000]	Year	B [mag]	ΔB [mag]	Tel.	Date [JD-2430000]	Year	B [mag]	ΔB [mag]	Tel.	Date [JD-2430000]	Year	B [mag]	ΔB [mag]	Tel.	Date [JD-2430000]	Year	B [mag]	ΔB [mag]	Tel.
5394.625	1955.7868	16.0	0.1	RA 122	6229.485	1958.0725	16.4	0.1	RL 60	7564.633	1961.7279	16.8	0.2	SA 50/40	8111.333	1963.2247	16.5	0.1	SB 120/80
5427.549	1955.8769	16.3	0.1	RA 122	6243.292	1958.1103	16.2	0.1	RL 60	7619.581	1961.8784	16.6	0.2	SA 50/40	8432.375	1964.1037	16.3	0.1	SB 120/80
5507.438	1956.0957	16.3	0.1	RA 122	6248.274	1958.1240	16.4	0.1	RL 60	7621.639	1961.8840	16.7	0.2	SA 50/40	9765.626	1967.7539	15.9	0.1	SA 90/65
5508.384	1956.0982	16.3	0.1	RA 122	6250.474	1958.1300	16.4	0.1	RL 60	7641.403	1961.9381	16.4	0.2	SA 50/40	9769.626	1967.7649	16.0	0.1	SA 90/65
5723.622	1956.6875	15.9	0.1	RA 122	6257.331	1958.1488	16.4	0.1	RL 60	7642.590	1961.9414	15.9	0.2	SA 50/40	9771.597	1967.7703	15.9	0.1	SA 90/65
5727.637	1956.6985	16.3	0.1	RA 122	6275.367	1958.1981	16.4	0.1	RL 60	7645.423	1961.9491	16.4	0.2	SA 50/40	9772.542	1967.7729	15.8	0.1	SA 90/65
5755.663	1956.7753	16.1	0.1	RA 122	6277.441	1958.2038	16.4	0.1	RL 60	7648.488	1961.9575	16.4	0.2	SA 50/40	9773.567	1967.7757	15.8	0.1	SA 90/65
5760.626	1956.7888	16.1	0.1	RA 122	6285.347	1958.2255	16.3	0.1	RL 60	7649.515	1961.9603	16.4	0.2	SA 50/40	9774.650	1967.7786	16.0	0.1	SA 90/65
5786.676	1956.8602	16.0	0.1	RA 122	6303.336	1958.2747	16.6	0.1	RL 60	7651.613	1961.9661	16.3	0.2	SA 50/40	9775.606	1967.7813	15.9	0.1	SA 90/65
5867.524	1957.0815	16.4	0.1	RL 60	6313.341	1958.3021	16.5	0.1	RL 60	7659.306	1961.9871	16.4	0.2	SA 50/40	9796.593	1967.8387	17.1	0.1	SA 90/65
5868.531	1957.0843	16.3	0.1	RA 122	6314.326	1958.3048	16.3	0.1	RL 60	7671.496	1962.0205	16.7	0.2	SA 50/40	9826.689	1967.9211	16.3	0.1	SA 90/65
5868.558	1957.0843	16.4	0.1	RL 60	6330.340	1958.3486	15.9	0.1	RL 60	7672.461	1962.0232	16.8	0.2	SA 50/40	9827.677	1967.9238	15.9	0.1	SA 90/65
5870.352	1957.0893	16.4	0.1	RL 60	6460.607	1958.7053	16.2	0.1	RL 60	7680.592	1962.0454	16.3	0.2	SA 50/40	9830.653	1967.9320	16.3	0.1	SA 90/65
5872.335	1957.0947	16.4	0.1	RL 60	6579.406	1959.0305	16.1	0.1	RL 60	7689.300	1962.0693	16.2	0.2	SA 50/40	9831.672	1967.9348	16.4	0.1	SA 90/65
5889.428	1957.1415	16.3	0.1	RL 60	6595.240	1959.0739	16.3	0.1	RL 60	7692.306	1962.0775	16.3	0.2	SA 50/40	9852.603	1967.9921	17.0	0.1	SA 90/65
5891.444	1957.1470	16.4	0.1	RL 60	6596.253	1959.0767	16.5	0.1	RL 60	7694.235	1962.0828	16.3	0.2	SA 50/40	9860.556	1968.0138	16.3	0.1	SA 90/65
5895.364	1957.1577	16.3	0.1	RL 60	6598.292	1959.0823	16.5	0.1	RL 60	7695.401	1962.0860	15.9	0.2	SA 50/40	9864.550	1968.0248	16.2	0.1	SA 90/65
5899.358	1957.1687	16.4	0.1	RL 60	6599.434	1959.0854	16.5	0.1	RL 60	7696.293	1962.0884	16.3	0.2	SA 50/40	9875.351	1968.0543	16.0	0.1	SA 90/65
5900.472	1957.1717	16.4	0.1	RL 60	6600.352	1959.0879	16.3	0.1	RL 60	7698.292	1962.0939	16.4	0.2	SA 50/40	9876.306	1968.0570	15.8	0.1	SA 90/65
5901.497	1957.1745	16.4	0.1	RL 60	6606.350	1959.1043	16.6	0.1	RL 60	7699.294	1962.0966	16.3	0.2	SA 50/40	9877.311	1968.0597	15.9	0.1	SA 90/65
5902.385	1957.1770	16.3	0.1	RL 60	6607.384	1959.1071	16.4	0.1	RL 60	7700.346	1962.0995	16.4	0.2	SA 50/40	9878.497	1968.0630	15.9	0.1	SA 90/65
5903.440	1957.1798	16.3	0.1	RL 60	6608.323	1959.1097	16.6	0.1	RL 60	7701.307	1962.1021	16.9	0.2	SA 50/40	9879.322	1968.0652	15.9	0.1	SA 90/65
5918.336	1957.2206	16.5	0.1	RL 60	6610.303	1959.1151	16.7	0.1	RL 60	7702.292	1962.1048	16.7	0.2	SA 50/40	9881.492	1968.0712	16.3	0.1	SA 90/65
5946.348	1957.2973	16.4	0.1	RL 60	6613.420	1959.1237	16.3	0.1	RL 60	7717.274	1962.1459	16.8	0.2	SA 50/40	9885.510	1968.0822	16.5	0.1	SA 90/65
6083.605	1957.6731	16.5	0.1	RL 60	6627.313	1959.1617	16.3	0.1	RL 60	7719.306	1962.1514	16.4	0.2	SA 50/40	9887.488	1968.0876	16.5	0.1	SA 90/65
6102.573	1957.7250	16.5	0.1	RL 60	6628.415	1959.1647	16.3	0.1	RL 60	7720.310	1962.1542	16.4	0.2	SA 50/40	9905.422	1968.1367	16.4	0.1	SA 90/65
6104.631	1957.7307	16.5	0.1	RL 60	6630.381	1959.1701	16.5	0.1	RL 60	7731.321	1962.1843	15.9	0.2	SA 50/40	9941.353	1968.2351	16.8	0.1	SA 90/65
6109.622	1957.7443	16.4	0.1	RL 60	6636.343	1959.1864	16.3	0.1	RL 60	7732.285	1962.1870	16.3	0.2	SA 50/40	9942.378	1968.2379	16.8	0.1	SA 90/65
6110.595	1957.7470	16.2	0.1	RL 60	6643.330	1959.2056	16.1	0.1	RL 60	7752.316	1962.2418	16.2	0.2	SA 50/40	9943.319	1968.2404	16.8	0.1	SA 90/65
6129.548	1957.7989	16.2	0.1	RL 60	6643.330	1959.2056	16.3	0.2	SA 50/40	7755.309	1962.2500	16.4	0.2	SA 50/40	9944.317	1968.2432	16.9	0.1	SA 90/65
6130.516	1957.8015	16.4	0.1	RL 60	6661.317	1959.2548	16.2	0.2	SA 50/40	7757.310	1962.2555	16.4	0.2	SA 50/40	9947.325	1968.2514	16.8	0.1	SA 90/65
6137.503	1957.8207	16.5	0.1	RL 60	6662.292	1959.2575	16.2	0.2	SA 50/40	7760.356	1962.2638	16.5	0.2	SA 50/40	10241.498	1969.0568	16.4	0.1	SA 90/65
6138.511	1957.8234	16.4	0.1	RL 60	6670.323	1959.2795	16.5	0.1	RL 60	7762.322	1962.2692	16.4	0.2	SA 50/40	10264.266	1969.1191	16.1	0.1	SA 90/65
6167.455	1957.9027	16.4	0.1	RL 60	6812.615	1959.6690	16.0	0.1	RL 60	7939.597	1962.7545	16.1	0.1	RA 122	10273.295	1969.1439	16.7	0.1	SA 90/65
6169.599	1957.9086	16.4	0.1	RL 60	6818.610	1959.6854	15.9	0.1	RL 60	7972.562	1962.8448	16.2	0.1	SB 120/80	10274.456	1969.1470	16.7	0.1	SA 90/65
6170.615	1957.9113	16.3	0.1	RA 122	6846.622	1959.7621	16.5	0.2	SA 50/40	7974.564	1962.8503	16.3	0.1	SB 120/80	10250.663	1969.1773	16.4	0.1	SA 90/65
6174.608	1957.9223	16.5	0.1	RL 60	6952.351	1960.0516	16.3	0.2	SA 50/40	7991.583	1962.8969	16.3	0.2	SA 50/40	10510.643	1969.7937	16.3	0.1	SA 90/65
6191.469	1957.9684	16.4	0.1	RL 60	7016.400	1960.2270	16.3	0.2	SA 50/40	7993.556	1962.9023	16.4	0.2	SA 50/40	10566.520	1969.9467	16.9	0.1	SA 90/65
6194.495	1957.9767	16.2	0.1	RL 60	7259.499	1960.8925	16.4	0.2	SA 50/40	7998.497	1962.9158	16.0	0.2	SA 50/40	10566.571	1969.9468	16.9	0.1	SA 90/65
6195.392	1957.9792	16.3	0.1	RL 60	7319.515	1961.0569	16.2	0.2	SA 50/40	8001.487	1962.9240	16.5	0.2	SA 50/40	10569.530	1969.9459	17.1	0.1	SA 90/65
6198.468	1957.9876	16.4	0.1	RL 60	7336.387	1961.1030	16.5	0.2	SA 50/40	8003.556	1962.9297	16.8	0.2	SA 50/40	10588.493	1970.0068	16.0	0.1	SA 90/65
6212.279	1958.0254	16.7	0.1	RL 60	7341.333	1961.1166	16.6	0.2	SA 50/40	8020.522	1962.9761	16.4	0.2	SA 50/40	10589.562	1970.0098	16.0	0.1	SA 90/65
6214.287	1958.0309	16.5	0.1	RL 60	7345.272	1961.1274	16.5	0.2	SA 50/40	8046.310	1963.0467	16.5	0.1	RA 122	10621.527	1970.0973	16.5	0.1	SA 90/65
6219.353	1958.0448	16.5	0.1	RL 60	7561.595	1961.7196	16.8	0.2	SA 50/40	8049.480	1963.0554	16.4	0.1	SB 120/80	10626.344	1970.1105	16.8	0.1	SA 90/65
6225.316	1958.0611	16.4	0.1	RL 60	7562.594	1961.7224	16.5	0.2	SA 50/40	8050.266	1963.0575	16.3	0.1	RA 122	10647.369	1970.1680	16.0	0.1	SA 90/65
6228.369	1958.0695	16.4	0.1	RL 60	7563.626	1961.7252	16.8	0.2	SA 50/40	8057.410	1963.0771	16.4	0.1	RA 122					
Date [JD-2430000]	Year	I [mag]	ΔI [mag]	Tel.	Date [JD-2430000]	Year	I [mag]	ΔI [mag]	Tel.	Date [JD-2430000]	Year	I [mag]	ΔI [mag]	Tel.	Date [JD-2430000]	Year	I [mag]	ΔI [mag]	Tel.
5755.645	1956.7752	13.9	0.1	RA 122	7364.304	1961.1795	13.9	0.2	SA 50/40	8050.281	1963.0576	14.6	0.1	RA 122	9881.506	1968.0712	13.9	0.1	SA 90/65
5815.572	1956.9393	14.0	0.1	RA 122	7365.352	1961.1823	13.9	0.2	SA 50/40	8057.392	1963.0770	14.6	0.1	RA 122	9885.325	1968.0822	14.3	0.1	SA 90/65
6101.603	1957.7224	13.9	0.1	RL 60	7366.372	1961.1851	13.8	0.2	SA 50/40	8799.463	1965.1087	14.0	0.2	SA 50/40	9887.503	1968.0876	14.4	0.1	SA 90/65
6109.602	1957.7443	13.9	0.1	RL 60	7367.306	1961.1877	13.9	0.2	SA 50/40	9763.640	1967.7485	13.9	0.1	SA 90/65	9905.462	1968.1368	14.0	0.1	SA 90/65
6130.538	1957.8016	14.0	0.1	RL 60	7368.326	1961.1905	13.8	0.2	SA 50/40	9765.647	1967.7540	14.1	0.1	SA 90/65	9941.338	1968.2350	14.8	0.1	SA 90/65
6138.535	1957.8235	13.9	0.1	RL 60	7369.309	1961.1932	14.0	0.2	SA 50/40	9769.651	1967.7650	13.9	0.1	SA 90/65	9942.358	1968.2378	14.8	0.1	SA 90/65
6170.507	1957.9110	14.1	0.1	RL 60	7370.317	1961.1959	13.8	0.2	SA 50/40	9771.620									

plates (in which the magnitude of the variable star was much fainter than the limit-magnitude of the plate). Hence these latter plates (about ten in *B*) were not useful to set meaning upper limits.

The merit of the material used in this work lays in its uniformity. In the blue, an emulsion 103a-O without filter was used. In the photographic-infrared, an emulsion I-N hypersensitized with an RG5 filter was used. The I-N/RG 5 combination transmits the spectral range 6800-8800 Å, and has a transmission curve similar to the 7000-9000 Å band-pass of the Cousins I_c band. Moreover, a large fraction of the plates (about 60%, see Tab. 1 and Tab. 2) were obtained with the 0.60m Loiano reflector and the 0.90/0.65m Asiago Schmidt telescope, having the same focal length (210 cm) and almost the same focal ratio (f/3.50 and f/3.23 respectively).

The irregular variability of the studied stars required a high frequency in the plate collection. As a consequence, for V582 Mon (KH 15D), a series of long duration observations (except for the seasonal, weather and full-moon interruptions) are available, in the years when the star began to show its variability and during the first evolution phases. For the first years this is true especially for the blue plates; for the subsequent years also in the infrared ones.

The magnitudes of comparison stars were taken from the blue photoelectric and photographic sequences of Walker (1956), after removing stars that appeared variable afterwards. The calibrations curves were constructed using the best two plates among the five ones obtained with the Schmidt telescope of the Hamburg-Bergedorf Observatory. The infrared comparison sequence was assembled using stars having a known colour index $B - V$ (Walker 1956) by means of the transformation formulae given by Allen (1955), taking care to choose only the non-reddened stars from the colour-colour-diagram $U - B$ vs $B - V$.

The magnitudes of V582 Mon (KH-15D) were determined through visual estimates with respect to its closest comparison stars. A stereoscopic Zeiss microscope at low and variable magnification was used. This method allows a comparison of the measurements obtained with different telescopes. For example, in the plates obtained with the Loiano reflector, the star falls in a region affected by coma, however the eye can perform a good estimation when comparison stars of comparable coma are chosen. On the basis of the thousands of magnitude estimations done with the above mentioned instruments, we estimate the mean errors in the magnitude estimations are: ± 0.1 mag for the 0.6m Loiano reflector plates, $\pm 0.05/0.1$ mag for the 1.2m Asiago reflector plates, ± 0.2 and ± 0.1 mag for the two 0.50/0.40m and 0.90/0.65m Asiago Schmidt telescope plates respectively.

3 DATA ANALYSIS AND RESULTS

Part of the analysis of the plates taken around NGC 2264 was published in Maffei (1966a,b). The detailed study of the more than 200 variable stars identified in the area of NGC 2264 (about one square degree) is an ongoing project. At present, this effort has been devoted to the variable V582 Mon (KH-15D). The infrared plates obtained with the two Asiago Schmidt telescopes were made available to J. A.




Figure 2. Uppermost panel: periodogram plot of the blue light curve in the best sampled interval JD 2436102 (1957.725) to JD 2438111 (1963.225). Dotted line shows the 1% false alarm significance level, under the hypothesis of fluctuations dominated by Poisson statistics. The timescale corresponding to the significant peak is 48.5 ± 0.3 days, in agreement with the phase-plots findings. Second panel: auto-correlations of the blue light curves, calculated with the DACF and ZDACF. These show a weak broad peak at around 48-49 days. Third panel: the first order structure function (SF) of the data in the same period, the SF indicates a drop around a timescale of 48 days. Bottom panel: the phase dispersion minimization (PDM) curve of the same data. The significant drop in the PDM gives a period of 48.4 ± 0.2 days, in agreement with the previous results. The results obtained with the different statistical methods indicate a period equal to the actual value.

Johnson, who analyzed the data and published a first set of results in Johnson & Winn (2004).

As shown in Tables 1 and 2, the blue light curve is better sampled. However for the years 1967-70, the number of useful plates is nearly equal for the two colours, since after the positive results obtained with the first observations in




Figure 3. Periodogram plot of the blue light curve in the interval JD 2435394 (1955.787) to JD 2436198 (1957.988). Dotted line shows the 1% false alarm significance level, under the hypothesis of fluctuations dominated by Poisson statistics. The curve does not display any characteristic timescale or periodical feature. Taking into account the good data sampling in 1957 (see Table 2) and the results displayed in Fig. 2, this implies that the variability/periodicity in the blue began in the first days of 1958.




Figure 4. Upper panel: periodogram plot of the infrared light curve in the interval JD 2436101 (1957.722) to JD 2438057 (1963.077). Dotted line shows the 1% false alarm significance level, under the hypothesis of fluctuations dominated by Poisson statistics. No comparable indication of a periodicity is seen in the infrared light curve periodogram. Lower panel: periodogram plot of the infrared light curve in the period 1967-1970. In this interval, the characteristic periodicity (48.4 day) is clearly observed also in the infrared data.

the infrared, PM decided to take both blue and IR images for every available observing night.

In the blue, V582 Mon appears with a constant magnitude from 1955 to 1957 (Fig. 1). The first clear sign of a significant weakening (0.4 mag) was observed on January 8, 1958. In the infrared, the star appeared with constant magnitude in the subsequent 4 years, and a first sure weakening (0.5 mag) is observed on November 16, 1961 (Fig. 1).

3.1 Temporal Analysis

In addition to the classical phase analysis, the temporal analysis of the data was carried out using well tested and

robust techniques suitable for an unevenly sampled time series: the Lomb-Scargle periodogram (LSP), the first order structure function (SF), the discrete auto-correlation function (DACF), and the phase dispersion minimization (PDM) method. These methods allow to investigate quantitatively the statistical temporal structure of variability, the auto-correlations, and the existence of characteristic time scales and periodicity.

The SF is equivalent to a power spectrum calculated in the time domain. It provides information on the type of variability and the range of the characteristic time scales that contribute to the fluctuations (see, e.g. Rutman 1978; Simonetti, Cordes, & Heeschen 1985). Steep drops in the SF indicate small variance and provide the signature of possible characteristic time scales, while the slope in the log-log representation gives the value of the power law spectrum index, and thus the nature of the variability. The DACF allows to investigate the level of auto-correlation in unevenly sampled data sets (see, e.g. Edelson & Krolik 1988; Hufnagel & Bregman 1992) without any interpolation. The number of points per time bin can vary greatly in the DACF, but data bins by equal population can be built together with Montecarlo estimations for peaks and uncertainties as in the Fisher z -transformed DACF (ZDACF, Alexander 1997). The LSP is a technique analogous to the Fourier analysis for discrete unevenly sampled data trains, useful to detect the strength of harmonic components with a certain angular frequency $\omega = 2\pi f$ (see, e.g. Scargle 1982; Horne & Baliunas 1986). Finally, the PDM method (see, e.g. Stellingwerf 1978; Jurkevich 1971) seeks to minimize the variance of the data at constant phase with respect to the mean value of the light curve. The PDM method has no preference for a particular periodical shape, it incorporates all the data directly into the test statistic and it is thus well suited to small and randomly spaced samples. A value is statistically significant when the PDM drops near zero.

In Fig. 2 we plot the LSP, DACF and ZDACF, SF, and PDM for the blue light curve data in the best sampled interval JD 2436102 (1957.725) to JD 2438111 (1963.225). The timescale corresponding to the significant peak is 48.5 ± 0.3 days for the LSP. The DACF and ZDACF also show a weak (value ~ 0.3) peak for a time lag at around 48-49 days. Both the SF and the PDM show a drop around the same value.

The apparently stable periodicity of the fluctuation is first observed in 1958. The period is in good agreement with that found in recent years with well sampled data. On the other hand the blue light curve does not display any characteristic timescale or periodical feature before 1958 (Fig. 3), reinforcing the evidence for a beginning of the eclipses around the first months of 1958.

The temporal analysis of the infrared light curve for 1957-1963 does not show any significant feature above statistical fluctuations in the LSP, DACF, SF, and PDM plots (upper panel of Fig. 4), which indicates no corresponding variability and periodicity in the infrared in these years. We believe that the variability occurred only some years later (possibly about 4 years later) in the infrared based on the periodical behaviour in the IR displayed in the last interval of our light curve (lower panel of Fig. 4), and by the $B - I$ colour analysis in Par. 3.4.




Figure 5. Optical blue magnitude light curve of V582 Mon (KH 15D) as function of the phase ϕ from our 1955-1970 data. The spectroscopic binary period (48.38 days) and the epoch 2452352.26 reported by Herbst et al. (2002) are used. All the data points reported in the plot are real observations, without any data point duplication. The eclipses appears to start at the end of the 1950s. The periodic variability in the subsequent years up to the 1970 seems rather sinusoidal, with slow descendent and increasing stages and no plateaus.

3.2 Blue Light Curve

After the first weakening, the variability amplitude increased until about 1.2 mag from the end of 1967 to the beginning of 1970. Applying the phase periodical analysis, the maximum probability period is 48.5 days, very similar to the ephemeris outlined by Herbst et al. (2002). The phase light curve constructed with the observations from 1967 to 1970, adopting the ephemeris of Herbst et al. (2002), shows a marked regularity.

Figure 5 shows the optical blue magnitude light curve of V582 Mon (KH 15D) as a function of the phase ϕ for our data. All the data points reported in the plot are real observations. The eclipses appear clearly to start at the end of the 1950s. The periodic variability in the subsequent years up to the 1970 seems rather sinusoidal, with slow decreasing and increasing stages and no evident plateaus, contrary to more recent observations. A phase shift of the light curve minimum was found in the infrared also by Johnson & Winn (2004).

3.3 Infrared Light Curve

The same behaviour is displayed by the infrared magnitude light curve in Fig. 6. In the photographic infrared (similar




Figure 6. Infrared magnitude light curve of V582 Mon (KH 15D) as a function of the phase ϕ from our 1955-1970 data. The spectroscopic binary period (48.38 days) and the epoch 2452352.26 reported by Herbst et al. (2002) are used. All the data points reported in the plot are real observations, without any data point duplication. In this plot the eclipses seems possibly to start at the beginning of 1960s, and the periodic variability in the subsequent years appear approximatively sinusoidal, with slow descendent and increasing stages and no evident plateaus.

to the I_c band as mentioned above) after the first weakening the amplitude increases, beginning in 1968, by 1.2 mag. The minima are coincident with those observed in the blue. The phase-constructed light curve shows a flat minimum in the years 1967-1970, similar to that observed in 1995, however it represents a smaller fraction of the period (0.2) than in the more recent data (0.4). The observed maximum does not appear as a pronounced plateau, as seen in recent observations. In addition, the transition between minimum and maximum values is much slower (0.2 against 0.07 in phase units); the infrared light curve is better described by a sinusoidal function rather than by a step-like structure, as observed in 1996-2002.

The infrared maximum appears approximatively at the same level as in recent years. We do not see a clear weakening, as instead suggested recently in the infrared magnitudes measured in the *real I* Cousin band. We must however cautiously underline that the two photometric systems are not identical. On the other hand, the variability amplitude increased from 1 mag to 3 mag.

3.4 Colour Indexes

The suggested 4-year difference in starting times between the blue and the IR periodicity is supported by the analysis




Figure 7. Upper panel: the $B - I$ colour index in function of the time. The plot is clearly in agreement with an infrared colour excess in the period 1955-1963, implying that the beginning of the periodical eclipsing variability in the blue was not followed by the infrared flux that remains rather constant in those years. Lower panel: the scatter plot between the $B - I$ colour index and the B magnitude brightness shows the same behaviour. The linear regression trend lines are reported. The slope is steeper for the 1955-1963 data (blue/darker square symbols) with respect to the slope of the 1967-1970 data (azure/lighter diamond symbols) implying that the star becomes redder when fainter during the 1955-1963 epoch.

of the $B - I$ colour indexes. For the calculation of the the $B - I$ index, we selected only the data from the two bands which were coincident within an interval of 1 day. Figure 7 shows the behaviour of the colour index in time as well as the correlation with the blue brightness, which fact confirms and underlines our finding that the beginning of the infrared variability occurred some years later (possibly 4-years) after that in the blue. In fact, the $B - I$ index shows a colour excess and a variability for 1955-1963, while it tends to remain constant in subsequent years (Fig. 7 upper panel). Moreover, the scatter plot $B - I$ versus B shows a steeper trend for the 1955-1963 data, implying that the star became redder while fainter during these years. We point out that there are several parallel B and I observations within one day, to support this hypotheses. Only in the subsequent years (1967-1970) does the behaviour of the colour index not depend on the source brightness, implying that the star shows the same periodical behaviour in the blue and infrared bands.

4 DISCUSSION AND CONCLUSIONS

We have presented the results of a long-term series of observations of the variable star V582 Mon (KH 15D) in the cluster NGC 2264, performed in the blue and near-infrared

bands between October 13, 1955 and March 1, 1970. This set of homogeneous data in two colours is unique to our knowledge for this period and reveals the start of the eclipse phase of V582 Mon in the two colours.

The analysis of our data provides the following important results:

- (i) in the blue, the star showed the first magnitude variations at the beginning of 1958;
- (ii) in the near infrared, the first magnitude variation was observed roughly 4 years later;
- (iii) over the above 4 year period the star showed an infrared excess, at least around minima.
- (iv) the period did not change;
- (v) the shape of the light curve appears different from the present one, i.e. a sinusoidal function rather than a curve with flat minima and maxima, as in Kearns & Herbst (1998).

A possible model to explain these characteristics and their evolution would be the following. Until 1958, V582 Mon (KH 15D) was surrounded by an edge-on disk or gaseous torus, which did not cause variations in the brightness. Subsequently, thin dust, able to absorb the blue radiation (but not the IR radiation) began to form, distributed as a sort of “banana” as in the model proposed by Barge & Viton (2003), but with nearby, or extended thinner edges as indicated by the absence of a flat maximum. Afterwards the dust aggregated into larger particles, and the selective absorption became total absorption. Due to the increasing number and dimension of the dust grains, the observed minima deepen. At the same time, the “banana” dust cloud (or its denser part) contracted, creating wider, flattened minima and steeper variations of the brightness to reach the actual shape and values of the light curve (Winn et al. 2004). This suggests the presence of a body or agglomerate of dust and/or stones less extended but denser and with sharper edges.

These hypotheses imply that in the last 40 years we have observed the birth of a giant planet near the star, as that discovered (with other techniques) in other nearby stars (see, e.g. Butler 2003; Marcy & Butler 1998). More likely, we advance the notion of the formation of a blob of material, from which one or more planets could form, or that could dissolve or evolve in an unforeseeable way. This is supported by the fact that V582 Mon (KH 15D) is a pre-main sequence star, i.e. in formation. Our data does not exclude other models. In any case, any proposed model must take into account the non-variability prior to 1958, the progressive increase of the absorption, the interval of 4 years separating the beginning of the blue and infrared variability and above all, the transformation from a sinusoidal light curve to an actual step-like structure characterized by flat maxima and minima. We believe that the evolution of this unique and peculiar stellar object must be followed in the future with several different instruments, and in particular with the same photometric and spectroscopic techniques used in the past.

5 ACKNOWLEDGMENTS

The authors wish to thank the referee W. Herbst for very useful comments that helped to improve this work.

M. Busso (Perugia University) and W. J. Burger (INFN-Perugia) are gratefully acknowledged for their contribution in the final revision of the paper.

REFERENCES

- Alexander, T. 1997, in *Astronomical Time Series*, Eds. Maoz, Sternberg & Leibowitz, Dordrecht: Kluwer, p. 163
- Allen, C. W. 1955, *Astrophysical Quantities*, London: Athlone, 180
- Agol, E., Barth, A. J., Wolf, S., Charbonneau, D. 2004, *ApJ*, 600, 781
- Badalian, H. S. Erastova, L. K. 1972, *Soobshch. Byurakan Obs.* n. 44
- Barge, P. Viton, M. 2003, *ApJ*, 593, L117
- Bryden, G., Różyczka, M., Lin, D. N. C., Bodenheimer, P. 2000, *ApJ*, 540, 1091
- Butler, P., in *Placet and Life: the Emerging Science of Astrobiology*, Baross, J. Sullivan, W. eds., Cambridge Univ. Press
- Chiang, E. I. & Murray-Clay, R. A. 2004, *ApJ*, 607, 913
- Deming, D., Charbonneau, D., Harrington, J. 2004, *ApJ*, 601, L87
- Edelson, R. A., Krolik, J. H. 1988, *ApJ*, 333, 646
- Flaccomio, E., Micela, G., Sciortino, S., Favata, F., Corbally, C., Tomaney, A. 1999, *A&A*, 345, 521
- Grinin, V. P. Tambovtseva, L. V. 2002, *Astronomy Letters*, 28, 601
- Johnson, J. A. Winn, J. N. 2004, *AJ*, 127, 2344
- Johnson, J. A., Marcy, G. W., Hamilton, C. M., Herbst, W., Johns-Krull, C. M. 2004, *AJ*, 128, 1265
- Jurkevich, I. 1971, *Ap&SS*, 13, 154
- Hamilton, C. M., Herbst, W., Mundt, R., Bailer-Jones, C. A. L., Johns-Krull, C. M. 2003, *ApJ*, 591, L45
- Hamilton, C. M., Herbst, W., Shih, C., Ferro, A. J. 2001, *ApJ*, 554, L201
- Herbst, W. Hamilton, C. M. 2003, *ASP Conf. Ser.* 294: *Scientific Frontiers in Research on Extrasolar Planets*, 231
- Herbst, W., Hamilton, C.M., Vrba, F.J., Ibrahimov, M.A., Bailer-Jones, C.A.L., Mundt, R., Lamm, M., Mazeh, T., Webster, Z.T., Haisch, K.E., et al. 2002, *PASP*, 114, 1167
- Horne, J. H., Baliunas, S. L. 1986, *ApJ*, 302, 757
- Hufnagel, B. R., Bregman, J. N. 1992, *ApJ*, 386, 473
- Kearns, K. E. Herbst, W. 1998, *AJ*, 116, 261
- Kearns, K. E., Eaton, N. L., Herbst, W., Mazzurco, C. J. 1997, *AJ*, 114, 1098
- Kholopov, P. N. (editor) 1985, *General Catalogue of Variable Stars* 4th edition, vol.1,2, Moscow: NAUKA
- Maffei, P. 1967, *ApJ*, 147, 802
- Maffei, P. 1966a, *Mem. Soc. Astron. Italiana*, 37, 459
- Maffei, P. 1966b, *Mem. Soc. Astron. Italiana*, 37, 475
- Marcy, G.W., Butler, R.P. 1998, *ARA&A*, 36, 57
- Park, B., Sung, H., Bessell, M. S., & Kang, Y. H. 2000, *AJ*, 120, 894
- Rosino, L., Grubissich, C., Maffei, P. 1957, *Contributi Oss. Astrofis. di Asiago* n. 82
- Rutman, J. 1978, *Proc. IEEE*, 66, 1048
- Scargle, J. D. 1982, *ApJ*, 263, 835
- Simonetti, J. H., Cordes, J. M., & Heeschen, D. S. 1985, *ApJ*, 296, 46
- Stellingwerf, R. F. 1978, *ApJ*, 224, 953
- Tokunaga, A. T., Dahm, S., Gässler, W., Hayano, Y., Hayashi, M., Iye, M., Kanzawa, T., Kobayashi, N., Kamata, Y., Minowa, Y., et al. 2004, *ApJ*, 601, L91
- Walker, M. F. 1961, *ApJ*, 133, 438
- Walker, M. F. 1956, *ApJS*, 2, 356
- Winn, J. N., Holman, M. J., Johnson, J. A., Stanek, K. Z., Garnavich, P. M. 2004, *ApJ*, 603, L45
- Winn, J. N., Garnavich, P. M., Stanek, K. Z., Sasselov, D. D. 2003, *ApJ*, 593, L121